

AUTOIMMUNE PRESCRIPTION FORM

Phone: 866.413.3156 Fax: 877.834.1231 / faxes@repharmacy.com



Patient Information

Name:				Parent/Guardian (if applicable):	
DOB:	Height:	Weight:	Phone:		Email:
Address:			City, State, Zip:		
Male	Female	First Dose of IVIG:	YES	NO	Prior IG Brands Used:
Specific Adverse Reaction w/ Prior Brands:					
Allergies:					

Diagnosis

Autoimmune Encephalopathy [G04.81](#)

Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) [G61.81](#)

Dermatopolymyositis [M33.90](#)

Guillain-Barre Syndrome (GBS) [G61.0](#)

Inflammatory Neuropathies [G61.89](#)

Multifocal Motor Neuropathy [G61.82](#)

Myasthenia Gravis (MG) [G70.0](#)

Myasthenia Gravis w/Acute Exacerbation [G70.01](#)

Pemphigoid [L12.0](#)

Pemphigus [L10.9](#)

Polymyositis [M33.20](#)

Pediatric Autoimmune Neuropsychiatric Disorders

Associated with Streptococcal Infections (PANDAS) [D89.89](#)

Stiff Person Syndrome [G25.82](#)

Other: _____

Prescription Information

Product: Pharmacist to Determine Physician Branded _____	
Intravenous Immunoglobulin Recommended dose 1gm/kg – 2gm/kg Infuse: _____ gm/kg Infuse: IV daily over _____ days; repeat every _____ weeks; _____ months Other: _____	Subcutaneous Immunoglobulin Pharmacy to determine # of sites unless alt # of sites indicated here: _____ Infuse: _____ times; _____ weeks; _____ months Other: _____
Infusion Rate: Please select one and provide complete information Pharmacist to determine per manufacturer recommendations Start at _____ mL/hour then increase by _____ mL/hour every _____ minutes to maximum rate _____ mL/hour	
IV Access: Peripheral PICC Port Other: _____	
IV Maintenance (Flushing): Dispense quantity sufficient Sodium chloride 0.9% 10mL prefilled syringe: flush IV access device with sodium chloride 1-10mL to maintain line patency Heparin 100 units/mL 5mL prefilled syringe: flush with 100 units/mL 3-5 mL as needed to maintain line patency	
Pretreatment: Dispense quantity sufficient Acetaminophen 325mg tablet: 1-2 tablets by mouth 15-30 minutes before each infusion Diphenhydramine 25mg capsule: 1-2 capsules by mouth 15-30 minutes before each infusion Other: _____	
Ancillary Supplies: Dispense ancillary supplies and equipment needed to provide home infusion therapy Refills: _____	
Labs: Labs will not be drawn on weekends/holidays. Not for STAT labs. Labs to be drawn: _____ Frequency of labs: _____	
Adverse/Anaphylactic Reactions: <i>Anaphylaxis kit to be used in the event of anaphylactic reaction and includes the below components:</i> Diphenhydramine 25mg capsule (qty 2), Diphenhydramine 50mg/mL 1mL vial (qty 1), Epinephrine injection auto-injector 0.3mg (>30kg pt) or 0.15mg (<30kg pt) (qty 1 two-pack), Sodium chloride 0.9% 500mL bag (qty 1), and Sodium chloride 0.9% 10mL prefilled syringe (qty 4)	
Nursing Orders: Nurse to administer IVIG and ancillary medications per orders. Skilled nursing visits for education of SCIG administration (not applicable if independent with therapy)	

Physician Information

Physician:		Office Contact:	
Address:		City, State, Zip:	
Phone:	Fax:	License:	NPI:
Number of Drugs Prescribed: _____ (Prescription is void if number of drugs are not indicated. Regulation 16 CCR 1717.3)			

Important Notice: To be administered by a healthcare professional. This document is intended solely for the designated recipient and contains confidential, privileged, or legally protected information. If you are not the intended recipient, you are prohibited from disseminating, distributing, or copying this document. Please notify the sender immediately if you have received this document in error and promptly destroy it. By signing this form and using our services, you authorize RE Pharmacy and its employees to act as your designated agent for prior authorization in dealings with medical and prescription insurance companies.

Prescriber's Signature:	Date:
-------------------------	-------